

**Non-native turtles in a peri-urban park in northern Milan (Lombardy, Italy):
species diversity and population structure**

By

CLAUDIO FOGLINI, ROBERTA SALVI

SUPPLEMENTARY MATERIAL

Figure S1. *T. scripta* hybrids. Usually, they showed faded orange or red postorbital stripes (faded red in a; orange in b) following the yellow facial marking, carapace with traces of yellow markings (a, b), or plastron with dark blotch or oculion on some scutes (c) (Seidel et al., 1999). Picture b and c are taken from the same specimen.



Figure S2. Specimen of *Graptemys pseudogeographica* Gray, 1831.



Figure S3. Specimen of *Pseudemy cfr. concinna* Le Conte, 1830.



Table S4. Capture-Marking-Recapture data for all sampled areas (output from Noremark, JHE model for close population).

Site	Captured turtles (n)	Recaptured turtles (n)	Population estimate (95% C.I.)
Cinisello Lake	13	13	13 (13)
Suzzani Lake	64	27	84 (73-102)
Bresso Lake	74	27	106 (90-129)
Breda Ditch	19	11	21 (19-26)

Figure S5. Juveniles of *T. scripta* hybrid (a) and *T. s. elegans* (b).



Figure S6. Nest-digging slider turtle female observed in July 2016. Photography by M.R. Gelso.



Figure S7. Juvenile slider turtle (*T. scripta scripta* or hybrid) observed in July 2016. Photography by M. Ghislandi.

