

Estimating abundance of the Stripeless tree-frog *Hyla meridionalis* by means of replicated call counts

FEDERICO CROVETTO, SEBASTIANO SALVIDIO, ANDREA COSTA *

DISTAV, University of Genova, Corso Europa 26, I-16132 Genova, Italy.

* *Corresponding author.* Email: andrea-costa-@hotmail.it

SUPPLEMENTARY MATERIAL




Fig. S1. Map of the sampling sites in the Cinque Terre National Park (lower right corner): A) Levanto; B) Monterosso; C) Riomaggiore. Maps for the current figure are distributed under an Open Data Commons/Open Database License by the ©OpenStreetMap contributors.

site/survey	2017.1	2017.2	2017.3	2018.1	2018.2	2018.3	2018.4	
1	NA	NA	NA		1	1	0	0
2	0	3	2	2	3	3	4	
3	1	6	4	2	3	3	4	
4	0	0	1	0	0	0	0	
5	5	4	0	1	3	3	3	
6	5	4	4	NA	NA	NA	NA	
7	0	0	0	0	0	0	0	
8	4	3	2	1	2	2	1	
9	2	3	4	1	1	1	0	
10	2	4	4	4	3	3	2	
11	4	1	1	3	3	2	0	
12	4	1	2	2	1	0	1	
13	1	0	0	0	0	0	0	
14	4	3	4	3	2	2	3	
15	1	2	1	0	0	2	0	
16	5	3	4	4	5	4	2	
17	2	1	2	0	1	0	0	
18	1	1	2	3	3	3	2	
19	0	0	1	2	2	0	0	
20	0	0	0	0	0	0	0	
21	0	0	0	0	0	0	0	
22	0	0	1	0	0	1	1	
23	1	3	1	1	1	1	1	
24	2	3	2	3	3	2	3	

Table S1. Counts of males at each site for 2017 and 2018 surveys.