

**Identification of biologically active fractions in the dermal secretions of the genus
Bombina (Amphibia: Anura: Bombinatoridae)**

IRYNA UDOVYCHENKO¹, OLEKSANDRA OSKYRKO¹, OLEKSII MARUSHCHAK², TETIANA
HALENOVA¹, OLEKSII SAVCHUK¹

¹*Educational and Scientific Centre "Institute of Biology and Medicine" Taras Shevchenko
National University of Kyiv, Ukraine, 03127, Hlushkova Avenue, 2.*

²*I. I. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kyiv,
01030, B. Khmelnytskogo str. 15.*

SUPPLEMENTARY MATERIAL

Fig. S1. Coomassie stained SDS-PAGE gel of the crude skin secretions of studied amphibian species: 1 – *B. bombina*, 2 – *B. variegata*, 3 – hybrid of *B. Bombina* and *B. variegata*, M – molecular weight markers




Fig. S2. Coomassie stained SDS-PAGE gel of the fractions #1-7 collected in the result of fractionation of *B. variegata* crude skin secretions. M – molecular weight markers




Fig. S3. Coomassie stained SDS-PAGE gel of the fractions #1-6 collected in the result of fractionation of *B. bombina* crude skin secretions. M – molecular weight markers




Fig. S4. Coomassie stained SDS-PAGE gel of the fractions #1-8 collected in the result of fractionation of the hybrid *B. bombina* × *B. variegata* crude skin secretions. M – molecular weight markers

