

**Threats of the emerging pathogen *Batrachochytrium salamandrivorans*  
(*Bsal*) to Italian wild salamander populations**

LORENZO DONDERO<sup>1</sup>, GIORGIA ALLARIA<sup>1</sup>, GIACOMO ROSA<sup>1</sup>, ANDREA COSTA<sup>1</sup>,  
GENTILE FRANCESCO FICETOLA<sup>2</sup>, ROBERTO COGONI<sup>3</sup>, ELENA GRASSELLI<sup>1</sup>, SEBASTIANO  
SALVIDIO<sup>1\*</sup>

<sup>1</sup>*Department of Earth, Environment and Life Sciences (DISTAV), University of Genova,  
Corso Europa 26, I-16132 Genova, Italy*

<sup>2</sup>*Department of Environmental Science and Policy, University of Milan, Via Celoria 10,  
20133 Milan, Italy*

<sup>3</sup>*Unione Speleologica Cagliariitana, Via Scarlatti 11, I-09045 Quartu St Elena, Italy*

*\*Corresponding author. Email: [sebastiano.salvidio@unige.it](mailto:sebastiano.salvidio@unige.it)*

**SUPPLEMENTARY MATERIAL**



**Fig. S1.** A) *Speleomantes sarrabusensis* after swabbing (photo F.G. Ficetola); B) corpse of *Speleomantes sarrabusensis* (photo F.G. Ficetola); C) *Salamandra salamandra* displaying skin wounds, highlighted by red circles (photo M. Fahrbach); D) Dorsal view of *Salamandra salamandra* before swabbing (photo S. Salvidio).