

Diet patterns of water green frogs (*Pelophylax esculentus* complex) in mixed population systems in Serbia

KATARINA V. BREKA^{1,*}, MILAN PLEČAŠ¹, NIKOLA VESOVIĆ¹, KATARINA STOJANOVIĆ¹, BORIS DUDIĆ¹ AND SRĐAN Ž. STAMENKOVIĆ¹

¹ *Institute of Zoology, Faculty of Biology, University of Belgrade, Studentski Trg 16, 11000 Belgrade, Serbia*

SUPPLEMENTARY MATERIAL

Table S1. Significant differences between diet composition in three localities using ANOSIM pairwise comparisons with sequential Bonferroni correction (global $\alpha < 0.05$). Locality abbreviations are as in the text.

	DTD	STR	JRG
DTD		0.0318	0.0052
STR	0.0318		0.0189
JRG	0.0052	0.0189	

Table S2. Contribution of the prey categories to the overall dissimilarity between the three localities from the SIMPER analysis. Prey categories abbreviations are as in the text.

Taxon	Average dissimilarity	Contribution %	Cumulative contribution%
Col.	17.85	20.82	20.82
Hym.	15.14	17.65	38.48
Lep_l	10.44	12.18	50.65
Dip	7.18	8.37	59.03
Hem.	6.21	7.24	66.27
Ara.	5.60	6.53	72.8
Ort.	4.91	5.73	78.53
Gas.	3.95	4.60	83.13
Lep.	3.27	3.82	86.94
Vert	2.49	2.90	89.85
Dip_l	2.19	2.56	92.4
Eph.	1.61	1.88	94.28
Odo.	1.43	1.67	95.95
Myr.	0.97	1.14	97.08
Cli.	0.90	1.05	98.13
Col_l	0.80	0.94	99.07
Mal.	0.80	0.93	100