Fluorescence in amphibians and reptiles: new cases and insights

LUCAS M. BOTELHO[1], SUZANA E. MARTINS[2], GREGORY MELOCCO[3], LUÍS F.C TOLEDO[4], IVAN SAZIMA[1,5], EDELCIO MUSCAT[1]

[1]*Projeto Dacnis, Estrada do Rio Escuro, 4754, Sertão das Cotias, Ubatuba, São Paulo, 11680-000, Brazil.*

[2]*IPBio – Instituto de Pesquisas da Biodiversidade, Reserva Betary, 18330-000, Iporanga, São Paulo, Brazil.*

[3]*Departamento de Ciências Farmacêuticas (Toxicologia e Fitopatologia), Farmácia, USP, 05508-000, São Paulo, São Paulo, Brazil.*

[4]*Laboratório de História Natural de Anfíbios Brasileiros (LaHNAB), Departamento de Biologia Animal, Instituto de Biologia, Unicamp, 13083-970, Campinas, São Paulo, Brazil.*

[5]*Museu de Diversidade Biológica, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, São Paulo, 13083-863, Brazil.*

SUPPLEMENTARY MATERIAL

Figure S1. Adult *Bokermannohyla alvarengai* with flash (A) and UV light (B).



Figure S2. Amphibian species with ocular fluorescence: *Hylodes phyllodes* photographed with flash (A) and under UV light (B); *Hylodes asper* photographed with flash (C) and under UV light (D).



Figure S3. Juvenile pitviper species with tail tip fluorescence: *Bothrops jararaca* photographed with flash (A) and under UV light (B); *Bothrops jararacussu* photographed with flash (C) and under UV light (D).